

File name: Insert VETAZINE Cream  
Scale: The label is 100% of full size  
Date: V01, 1008

### **VETAZINE™ CREAM**

Modern Veterinary Therapeutics, LLC  
TRIAMCINOLONE ACETONIDE CREAM, USP 0.1%.  
For Topical Use on Dogs Only

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**DESCRIPTION:** VETAZINE™ Cream (triamcinolone acetonide) provides 1 mg triamcinolone acetonide per gram (0.1%) in a vanishing cream base containing propylene glycol, emulsifying wax, mineral oil, sorbitol, cetyl palmitate, sorbic acid, potassium sorbate and purified water.

**ACTIONS:** VETAZINE™ Cream contains a corticosteroid that provides prompt relief of itching, burning, inflamed skin lesions by virtue of its anti-inflammatory, antipruritic and anti-allergic effects.

**INDICATIONS:** VETAZINE™ Cream is indicated for topical treatment of allergic dermatitis and summer eczema in dogs.

**CONTRAINDICATIONS:** VETAZINE™ Cream should not be used ophthalmically.

**WARNINGS:** VETAZINE™ Cream is indicated for use on dogs only. Do not use VETAZINE™ Cream on animals which are raised for food production.

Absorption of triamcinolone acetonide through topical application on the skin and by licking does occur. Therefore, dogs receiving VETAZINE™ Cream therapy should be observed closely for signs of polydipsia, polyuria and increased weight gain, particularly when used over large areas or for extended periods of time.

Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection to animals may induce the first stage of parturition if used during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta and metritis.

Additionally, corticosteroids administered to dogs, rabbits and rodents during pregnancy have resulted in cleft palate in offspring. Corticosteroids administered to dogs during pregnancy have also resulted in other congenital anomalies including deformed forelegs, phocomelia and anasarca.

**PRECAUTIONS:** If local infection exists, suitable concomitant antimicrobial therapy should be administered. If favorable response does not occur promptly, application of VETAZINE™ Cream should be discontinued until the infection is adequately controlled by appropriate measures. Avoid ingestion. Oral or parenteral use of corticosteroids, depending on dose, duration and specific steroid, may result in inhibition of endogenous steroid production following drug withdrawal.

**SIDE EFFECTS:** SAP and SGPT (ALT) enzyme elevations, polydipsia and polyuria have occurred following parenteral or systemic use of synthetic corticosteroids in dogs. Vomiting and diarrhea (occasionally bloody) have been observed in dogs.

Cushing's Syndrome in dogs has been reported in association with prolonged or repeated steroid therapy.

**DOSAGE AND ADMINISTRATION:** Apply VETAZINE™ Cream by rubbing into the affected areas two to four times daily for 4 to 10 days.

**HOW SUPPLIED:** VETAZINE™ Cream is supplied in 28.4 & 85.2 gram tubes.

**STORAGE:** Store at room temperature; avoid freezing. Do not store above 30°C (86°F).

ANADA 200-459, Approved by FDA.

Manufactured for:  
Modern Veterinary Therapeutics, LLC  
Coral Gables, Florida 33146 - USA

Rev#1008

